

# Exceptional outcomes. Proven again.



Japan IDE Clinical Study Results: Proven patency in complex SFA lesions.

**88%** 12-month primary patency in long, complex SFA lesions (n = 103)<sup>1</sup>

- 21.8 cm average lesion length
- 65.7% chronic total occlusions (CTOs)
- 84.5% TASC II C&D lesions



## 12-Month Patencies: GORE® VIABAHN® Endoprosthesis Japan IDE Clinical Study\*



\* GORE® VIABAHN® Endoprosthesis Japan IDE Clinical Study demonstrated 12-month primary patency of 92% as defined by evidence of flow with no Target Lesion Revascularization (TLR). The same study demonstrated 88% 12-month primary patency when defined by PSVR of < 2.5 without a TLR.




# Proven Patency for Complex SFA Lesions.

**359** Limbs Studied

**255** CTOs

**21 cm** Average Lesion Length\*

**78%** Average Primary Patency\*\*



STUDY	NUMBER OF LIMBS	LESION LENGTH (cm)	CTOs (%)	12-MONTH PRIMARY PATENCY	12-MONTH SECONDARY PATENCY
Japan IDE Clinical Study <sup>1</sup>	103	22	66	<b>88</b>	<b>98</b>
Gore VIPER Clinical Study <sup>2</sup>	119	19	56	73	92
VIASTAR Trial <sup>3</sup>	66	19	79	78	90
25 cm Trial <sup>4</sup>	71	27	93	67	97
<b>Combined Results</b> (Weighted average, as appropriate)	<b>359</b>	<b>21</b>	<b>70</b>	<b>78</b>	<b>94</b>

Read the Japan IDE Clinical Study Abstract at [goremedical.com/ap/viabahn](http://goremedical.com/ap/viabahn)

\* Weighted Average Lesion Length

\*\* 12-Month Weighted Average Primary Patency

1. Ohki T, Kichikawa K, Yokoi H, *et al.* Outcomes of the Japanese multicenter Viabahn trial of endovascular stent grafting for superficial femoral artery lesions. *Journal of Vascular Surgery* 2017;66(1):130-142.e1.
2. Saxon RR, Chervu A, Jones PA, *et al.* Heparin-bonded, expanded polytetrafluoroethylene-lined stent graft in the treatment of femoropopliteal artery disease: 1-year results of the VIPER (Viabahn Endoprosthesis with Heparin Bioactive Surface in the Treatment of Superficial Femoral Artery Obstructive Disease) Trial. *Journal of Vascular & Interventional Radiology* 2013;24(2):165-173.
3. Lammer J, Zeller T, Hausegger KA, *et al.* Heparin-bonded covered stents versus bare-metal stents for complex femoropopliteal artery lesions: the randomized VIASTAR trial (Viabahn endoprosthesis with PROPATEN bioactive surface [VIA] versus bare nitinol stent in the treatment of long lesions in superficial femoral artery occlusive disease). *Journal of the American College of Cardiology* 2013;62(15):1320-1327.
4. Zeller T, Peeters P, Bosiers M, *et al.* Heparin-bonded stent-graft for the treatment of TASC II C and D femoropopliteal lesions: the Viabahn-25 cm Trial. *Journal of Endovascular Therapy* 2014;21(6):765-774.



**W. L. GORE & ASSOCIATES, INC.**  
Flagstaff, AZ 86004

+65.67332882 (Asia Pacific)  
00800.6334.4673 (Europe)  
800.437.8181 (United States)  
928.779.2771 (United States)

[goremedical.com](http://goremedical.com)

Products listed may not be available in all markets.

GORE®, VIABAHN®, and designs are trademarks of W. L. Gore & Associates. © 2017 W. L. Gore & Associates, Inc. AW1471-EN1 NOVEMBER 2017

Consult Instructions for Use