

GORE® PRECLUDE® Pericardial Membrane is demonstrated to have low risk of infection¹⁻¹⁴

0.7% Low risk of deep infection in patients implanted with GORE® PRECLUDE® Pericardial Membrane¹⁻¹⁴

Patients implanted with GORE® PRECLUDE® Pericardial Membrane

N = 2855

Patients with deep infection

19 (0.7%)

Left ventricular assist device (LVAD) and total artificial heart (TAH) implantations increase the risk of infection, regardless of utilizing GORE® PRECLUDE® Pericardial Membrane during surgery.¹⁵⁻¹⁶

6.4% - 53.7%¹⁷

Meta analysis: Infection rate range of LVAD and TAH surgeries

7.7% Clinical studies / reports utilizing GORE® PRECLUDE® Pericardial Membrane have shown an infection rate within the range associated with LVAD and TAH surgeries.^{9-11, 18-19}

Reoperation patients implanted with an LVAD or TAH and GORE® PRECLUDE® Pericardial Membrane


N = 65

Patients with deep infection

5 (7.7%)

“There were no perioperative complications or deaths attributable to the membrane [GORE® PRECLUDE® Pericardial Membrane]. There were no membrane [GORE® PRECLUDE® Pericardial Membrane]-related infections”¹⁰

Reprinted from *Operative Techniques in Thoracic and Cardiovascular Surgery*, Vol:19, Osami Honjo, Vivek Rao, Implantation of HeartWare Left Ventricular Assist Device in Pediatric Population, Figure 8a, page 93, 2014, with permission from Elsevier. Image courtesy Osami Honjo, M.D., Ph.D. Used with permission.



For more information on GORE® PRECLUDE® Pericardial Membrane clinical study data contact your clinical sales associate.



References

1. Mestres C-A, Comas JV, Ninot S, *et al.* The use of polyetherurethane urea (Mitrathane) and polytetrafluoroethylene (Gore-Tex) membranes for pericardial closure: initial clinical results. *Thai Journal of Surgery* 1986;7(4):125-128.
2. Minale C, Nikol S, Hollweg G, Mittermayer C, Messmer BJ. Clinical experience with expanded polytetrafluoroethylene Gore-Tex surgical membrane for pericardial closure: a study of 110 cases. *Journal of Cardiac Surgery* 1988;3(3):193-201.
3. Amato JJ, Cotroneo JV, Galdieri RJ, Alboliras E, Antillon J, Vogel RL. Experience with the polytetrafluoroethylene surgical membrane for pericardial closure in operations for congenital cardiac defects. *Journal of Thoracic & Cardiovascular Surgery* 1989;97(6):929-934.
4. Loebe M, Alexi-Meskishvili V, Weng Y, Hausdorf G, Hetzer R. Use of polytetrafluoroethylene surgical membrane as a pericardial substitute. *Texas Heart Institute Journal* 1993;20(3):213-217.
5. Alexi-Meskishvili V, Weng Y, Uhlemann F, Lange PE, Hetzer R. Prolonged open sternotomy after pediatric open heart operation: experience with 113 patients. *Annals of Thoracic Surgery* 1995;59(2):379-383.
6. Jacobs JP, Iyer RS, Weston JS, *et al.* Expanded PTFE membrane to prevent cardiac injury during resternotomy for congenital heart disease. *Annals of Thoracic Surgery* 1996;62(6):1778-1782.
7. Minale C, Hollweg G, Nikol S, Mittermayer CH, Messmer BJ. Closure of the pericardium using expanded polytetrafluoroethylene GORE-TEX surgical membrane: clinical experience. *Thoracic & Cardiovascular Surgeon* 1987;35(5):312-315.
8. Revuelta JM, Garcia-Rinaldi R, Johnston Jr. RH, Vaughan III. GD. Implantation of pericardial substitutes. *Annals of Thoracic Surgery* 1985;39(2):190-191.
9. Leprince P, Rahmati M, Bonnet N, *et al.* Expanded polytetrafluoroethylene membranes to wrap surfaces of circulatory support devices in patients undergoing bridge to heart transplantation. *European Journal of Cardio-thoracic Surgery* 2001;19(3):302-306.
10. Vitali E, Russo C, Tiziano C, Lanfranconi M, Bruschi G. Modified pericardial closure technique in patients with ventricular assist device. *Annals of Thoracic Surgery* 2000;69(4):1278-1279.
11. Copeland JG, Arabia FA, Smith RG, Covington D. Synthetic membrane neo-pericardium facilitates total artificial heart explantation. *Journal of Heart & Lung Transplantation* 2001;20(6):654-656.
12. Ozeren M, Han U, Mavioglu I, *et al.* Consequences of PTFE membrane used for prevention of re-entry injuries in rheumatic valve disease. *Cardiovascular Surgery* 2002;10(5):489-493.
13. Kajiwarra H, Ichikawa Y, Yamazaki I, Uranaka Y, Hamada T. Experience with expanded polytetrafluoroethylene surgical membrane (Gore-Tex surgical membrane) for coronary artery bypass grafting does Gore-Tex surgical membrane induce postoperative mediastinitis? *ASAIO Journal* 2003;49(2):155.
14. Bezon E, Maguid YA, Gueret G, Choplain JN, Aziz AA, Barra JA. Wrapping of the left internal thoracic artery with an expanded polytetrafluoroethylene membrane. *Annals of Thoracic Surgery* 2006;8(1):383-385.
15. Holman WL, Skinner JL, Waites KB, Benza RL, McGiffin DC, Kirklin JK. Infection during circulatory support with ventricular assist devices. *Annals of Thoracic Surgery* 1999;68(2):711-716.
16. Goldstein DJ, Oz MC, Rose EA. Implantable left ventricular assist devices. *New England Journal of Medicine* 1998;339(21):1522-1533.
17. Maniar S, Kondareddy S, Topkara VK. Left ventricular assist device-related infections: past, present and future. *Expert Review of Medical Devices* 2011;8(5):627-634.
18. Holman WL, Bourge RC, Zorn GL, Brantley LH, Kirklin JK. Use of expanded polytetrafluoroethylene pericardial substitute with ventricular assist devices. *Annals of Thoracic Surgery* 1993;55(1):181-183.
19. Kornberger A, Walter V, Khalil M, *et al.* Suspected involvement of EPTFE membrane in sterile intrathoracic abscess and pericardial empyema in a multi-allergic LVAD recipient: a case report. *Journal of Cardiothoracic Surgery* 2015;10(1):99.



W. L. GORE & ASSOCIATES, INC.
Flagstaff, AZ 86004

+65 67332882 (Asia Pacific) 800 437 8181 (United States)
00800 6334 4673 (Europe) 928 779 2771 (United States)

goremedical.com

 Consult Instructions for Use

INDICATIONS FOR USE: Reconstruction or repair of the pericardium. **CONTRAINDICATIONS:** Not for reconstruction of CARDIOVASCULAR DEFECTS such as cardiac, great vessel, and peripheral vascular, DURA MATER, HERNIAS. Use of this product in applications other than those indicated has the potential for serious complications, such as suture pullout or failure of the repair (aneurysm formation). Refer to *Instructions for Use* at goremedical.com for a complete description of all warnings, precautions and contraindications. Rx Only

Products listed may not be available in all markets.

GORE, PRECLUDE and designs are trademarks of W. L. Gore & Associates.
© 2019 W. L. Gore & Associates, Inc. AY1586-EN1 NOVEMBER 2019